

Correction to “Atmospheric temperature change detection with GPS radio occultation 1995 to 2008”

A. K. Steiner, G. Kirchengast, B. C. Lackner, B. Pirscher, M. Borsche, and U. Foelsche

Received 8 January 2010; published 13 February 2010.

Citation: Steiner, A. K., G. Kirchengast, B. C. Lackner, B. Pirscher, M. Borsche, and U. Foelsche (2010), Correction to “Atmospheric temperature change detection with GPS radio occultation 1995 to 2008,” *Geophys. Res. Lett.*, 37, L03704, doi:10.1029/2010GL042427.

[1] In the paper “Atmospheric temperature change detection with GPS radio occultation 1995 to 2008” by A. K. Steiner et al. (*Geophys. Res. Lett.*, 36, L18702, doi:10.1029/2009GL039777), Figure S7 of the auxiliary material is incorrect and the corrected version appears here.¹

Auxiliary material for Paper 2010GL042427

Correction to "Atmospheric temperature change detection with GPS radio occultation 1995 to 2008"

A. K. Steiner, G. Kirchengast, B. C. Lackner, B. Pirscher, M. Borsche, and U. Foelsche

The auxiliary material contains the corrected version of Figure S7.




Figure S7 corrected: Trend pattern (temperature change in [K/decade]) of RO monthly mean temperature anomalies at 10° zonal resolution (**a**, left) compared to corresponding HadAT2 radiosonde temperature anomalies (**a**, right) for the GPS/Met plus CHAMP period (1995, 1997, 2001–2008) and correspondingly for the CHAMP period only (2001–2008) (**b**). Also shown, for context, is the trend pattern of HadAT2 radiosonde temperatures for the continuous period 1995–2008 at 10° zonal resolution (**c**, left) and at study resolution (50S–20S, 20S–20N, 20N–50N) (**c**, right).